

Recent Results from KamLAND

Koichi Ichimura for the KamLAND Collaboration
University of California, Berkeley, California 94720, USA

The main goal of the KamLAND reactor $\bar{\nu}_e$ experiment is a search for $\bar{\nu}_e$ oscillation using inverse- β decay reaction in 1,000 ton of ultra-pure liquid scintillator. The data-set is 1490.8 days from Mar. 2002 to May 2007. The best-fit oscillation parameters are $\Delta m_{21}^2 = 7.58_{-0.13}^{+0.14}$ (stat.) ± 0.15 (syst.) $\times 10^{-5}$ eV² and $\tan^2 \theta_{12} = 0.56_{-0.07}^{+0.10}$ (stat.) $_{-0.06}^{+0.10}$ (syst.). The statistical significance for reactor $\bar{\nu}_e$ disappearance is 8.8σ , and an undistorted $\bar{\nu}_e$ energy spectrum is disfavored at $> 5\sigma$.

1. Introduction

KamLAND(**K**amioka **L**iquid scintillator **A**nti-**N**eutrino **D**etector) investigates neutrino oscillation parameters by observing electron anti-neutrinos ($\bar{\nu}_e$) from distant nuclear reactors. Previously, KamLAND reported the first evidence of $\bar{\nu}_e$ disappearance [1] and evidence of spectral distortion of the reactor $\bar{\nu}_e$ energy spectrum [2]. Furthermore, KamLAND published the first investigation of geologically produced anti-neutrinos (geo-neutrinos) from radioactive decay in the Earth[3].

In our recent analysis [4], we have enlarged the fiducial volume radius from 5.5m to 6.0m, extended the analysis threshold down to the inverse- β decay energy threshold and reduced the systematic uncertainties in the number of target protons and the background.

2. Detection method and analysis improvements

2.1. Detection method

$\bar{\nu}_e$ are detected via the inverse β decay reaction, $\bar{\nu}_e + p \rightarrow n + e^+$. The emitted positron annihilates with an electron and emits two 0.511MeV γ -rays. This signal appears as a prompt signal and its energy consist of the positron kinetic energy and these two γ -rays. The prompt scintillation light from the e^+ gives a measure of the $\bar{\nu}_e$ energy, $E_{\bar{\nu}_e} \simeq E_{prompt} + \bar{E}_n + 0.8\text{MeV}$, where $\bar{E}_n$ is the average neutron recoil energy, O(10keV). The neutron thermalizes by elastic scattering in the liquid scintillator and captures on a proton after $207.5 \pm 2.8\mu\text{sec}$, creating a deuteron with 2.2MeV γ emission. The neutron capture is the delayed signal. By using the delayed coincidence method, the backgrounds that do not have timing or space correlations are significantly reduced.

2.2. Off-axis calibration

We recently used an off-axis calibration system to investigate the vertex position reconstruction performance. Up to 5.5m radius, the vertex reconstruction deviation is within 3cm at any (R, θ, ϕ) , corresponding to 1.6% fiducial volume uncertainty. $^{12}\text{B}/^{12}\text{N}$ spallation products are used to extrapolate this uncertainty up to 6.0m fiducial radius, resulting in 1.8% uncertainty. The vertex and energy resolution are also estimated from off-axis and z-axis calibration data. The estimated vertex resolution is $\sim 12\text{cm}/\sqrt{E(\text{MeV})}$ and the energy resolution is $6.5\%/\sqrt{E(\text{MeV})}$.

2.3. Likelihood selection

In Ref.[2], 5.5m fiducial radius and 2.6MeV prompt energy threshold were applied to the reactor $\bar{\nu}_e$ analysis. To get more statistics and to use the full reactor energy spectrum, a larger fiducial volume and a lower energy threshold are required. However, accidental coincidences due to ^{40}K , ^{210}Bi and ^{208}Tl increase near the balloon

surface ($R = 6.5\text{m}$) reducing the signal-noise ratio. To maintain high efficiency for $\bar{\nu}_e$ detection, we apply a likelihood selection using event characteristics. Firstly we construct a probability density function (PDF) for accidental coincidence events, $f_{acci}(E_p, E_d, \Delta R, \Delta T, R_p, R_d)$ by using an off-time (10ms to 20 sec) window. A PDF for the $\bar{\nu}_e$, $f_{\bar{\nu}_e}(E_p, E_d, \Delta R, \Delta T, R_p, R_d)$ is constructed from a MC simulation. For the E_{prompt} distribution, we select an no-oscillation reactor spectrum including a contribution from geo-neutrinos. A discriminator value, $L = \frac{f_{\bar{\nu}_e}}{f_{acci} + f_{\bar{\nu}_e}}$ is calculated for all candidate pairs that pass the primary cuts. We choose a selection value L_i^{cut} in E_p bins of 0.1 MeV, where L_i^{cut} is the value which has the maximum figure-of-merit. Table I shows the summary of the systematic uncertainties in the analysis. The dominant uncertainties come from the reactor-related ones.

Table I: Estimated systematic uncertainties relevant for the neutrino oscillation parameters.

	Detector-related(%)	Reactor-related(%)		
Δm_{21}^2	Energy scale	1.9	$\bar{\nu}_e$ spectra	0.6
Event rate	Fiducial Volume	1.8	$\bar{\nu}_e$ spectra	2.4
	Energy threshold	1.5	Reactor power	2.1
	Efficiency	0.6	Fuel composition	1.0
	Cross section	0.2	Long-lived nuclei	0.3

2.4. Background

The dominant background is caused by the $^{13}\text{C}(\alpha, n)^{16}\text{O}$ reaction. The primary α source is the decay of ^{210}Po , a decay-daughter of ^{222}Rn introduced during the liquid scintillator filling. In Ref.[2], a 32% error for the reaction to ^{16}O ground state was assigned from uncertainties of ^{210}Po rate, the cross section to ^{16}O ground state, neutron angular distribution and proton quenching. A conservative 100% error was assigned to the excited states since the cross section was not well known. To measure the (α, n) prompt spectrum, a detector calibration with a $^{210}\text{Po}^{13}\text{C}$ source was performed. The expected spectrum was also simulated by the $^{13}\text{C}(\alpha, n)^{16}\text{O}$ cross sections from Ref.[5][6] and compared with the data. The new cross section measurement[5] has a 4% uncertainty in the total cross section, but it does not specify the final state separately. Thus the subtraction of the reaction rate of the excited states estimated from the cross section in Ref.[6] is needed for this new measurement. The first ^{16}O excited state emits e^+e^- , but they cannot escape from the source capsule. In KamLAND, these events are observed at $\sim 1.02\text{MeV}$ from the annihilation gammas. Therefore, the reaction rate of each state can be measured separately by $^{210}\text{Po}^{13}\text{C}$ calibration source. The rate of the second excited state was consistent with the calculation[6] without scaling. For the first excited state, the calculated spectrum agreed with the calibration data if scaled by 0.6. After subtraction of the excited state cross sections, the ground state was obtained. The reaction rate of the ground state had to be scaled by 1.05 to get the calculated ground state to be in good agreement with the data. With these improvements, we assign an uncertainty of 11% for the ground state and 20% for the excited states. Finally, the number of background events due to $^{13}\text{C}(\alpha, n)^{16}\text{O}$ is estimated to be 182 ± 21.7 events. Table II shows the summary of the backgrounds.

3. Results

We observe 1609 events in our data-set. With no $\bar{\nu}_e$ disappearance, we expect 2179 ± 89 events from the reactors. The background in this analysis is 276.1 ± 23.5 events excluding the geo-neutrino signal. Figure 1 shows the prompt energy spectrum of selected $\bar{\nu}_e$ events and the fitted backgrounds. The unbinned data is assessed with a maximum likelihood fit to two-flavor neutrino oscillation, simultaneously fitting the geo-neutrino contribution. The method uses the absolute time of the event and accounts for time variations in the reactor flux. Earth matter oscillation effects are also included.

Table II: Summary of estimated backgrounds

Background	contribution
Accidentals	80.5 ± 0.1
$^8\text{He}/^9\text{Li}$	13.6 ± 1.0
Fast neutrons and Atmospheric ν	< 9.0
$^{13}\text{C}(\alpha, n)^{16}\text{O}$ G.S.	157.2 ± 17.3
$^{13}\text{C}(\alpha, n)^{16}\text{O}$ $^{12}\text{C}(n, n\gamma)^{12}\text{C}$ (4.4 MeV γ)	6.1 ± 0.7
$^{13}\text{C}(\alpha, n)^{16}\text{O}$ 1st excited state (6.05 MeV e^+e^-)	15.2 ± 3.5
$^{13}\text{C}(\alpha, n)^{16}\text{O}$ 2nd excited state (6.13 MeV γ)	3.5 ± 0.2
Total	276.1 ± 23.5




Figure 1: Prompt energy spectrum of $\bar{\nu}_e$ candidate events. All histograms corresponding to reactor spectra and expected backgrounds incorporate the energy-dependent selection efficiency (top). The shaded background and geo-neutrino histograms are cumulative. Statistical uncertainties are shown for the data. The band on the blue histogram indicates the event rate systematic uncertainty.

The best-fit is shown in Figure 1. The best-fit oscillation parameters are $\Delta m_{21}^2 = 7.58_{-0.13}^{+0.14} (stat.) \pm 0.15 (syst.) \times 10^{-5} \text{ eV}^2$ and $\tan^2 \theta_{12} = 0.56_{-0.07}^{+0.10} (stat.) \pm 0.06_{-0.06}^{+0.10} (syst.)$. A scaled reactor spectrum with no distortion from neutrino oscillation is excluded at more than 5σ . Figure 2 shows the allowed contours in the neutrino oscillation parameter space. Only the so-called LMA-I region remains and the LMA-II and LMA-0 regions are disfavored at $> 4\sigma$. A combined oscillation analysis of KamLAND and other solar neutrino experiments under the assumption of CPT invariance gives $\Delta m_{21}^2 = 7.59 \pm 0.21 \times 10^{-5} \text{ eV}^2$ and $\tan^2 \theta_{12} = 0.47_{-0.05}^{+0.06}$. The Δm_{21}^2 parameter is strongly determined by the KamLAND experiment.

The L_0 / E distribution is suitable for illustrating the oscillatory behavior, see Figure 3. In this figure, the data and the best-fit spectrum are divided by the expected spectrum without oscillation. L_0 is the flux-weighted effective baseline and is 180 km. Two oscillation periods are seen, the KamLAND data is in excellent agreement with the oscillation scenario.

To determine the number of geo-neutrinos, we fit the normalization of the $\bar{\nu}_e$ energy spectrum from U and Th decay chains simultaneously with the neutrino oscillation parameters using the KamLAND and solar data. Fixing the Th/U mass ratio from planetary data to 3.9, the best-fit value is 73 ± 27 events.

4. Conclusion

The KamLAND experiment has determined a precise value for the neutrino oscillation parameter Δm_{21}^2 and stringent constraints on θ_{12} using reactor anti-neutrinos from 55 Japanese reactors over a five years period. The observed neutrino spectrum is distorted and the significance of the spectral distortion is $> 5\sigma$ and no-spectral distortion is strongly disfavored. A two-neutrino oscillation analysis of the KamLAND data with 0.9 MeV threshold gives $\tan^2 \theta_{12} = 0.56_{-0.07}^{+0.10} (stat.) \pm 0.06_{-0.06}^{+0.10} (syst.)$ and $\Delta m_{21}^2 = 7.58_{-0.13}^{+0.14} (stat.) \pm 0.15 (syst.) \times 10^{-5} \text{ eV}^2$. The previous result [2] was $\tan^2 \theta_{12} = 0.46_{-0.13}^{+0.23}$ and $\Delta m_{21}^2 = 7.9_{-0.3}^{+0.4} \times 10^{-5} \text{ eV}^2$. The uncertainties of neutrino oscillation parameters are significantly reduced and the systematic and statistical uncertainty are now comparable. The LMA-II and LMA-0 which were both allowed in Ref. [2] are now disfavored at $> 4\sigma$ and only LMA-I remains. These results are also published in Ref. [4].




Figure 2: Allowed region for neutrino oscillation parameters from KamLAND and solar neutrino experiments. The side panels show the $\Delta\chi^2$ -profiles for KamLAND(dashed),solar experiments(dotted) and the combination of the two(solid).




Figure 3: Ratio of the background subtracted $\bar{\nu}_e$ spectrum to the expectation without oscillation as a function of L_0/E . L_0 is the effective baseline taken as a flux weighted average ($L_0 = 180$ km). The energy bins are equal probability bins of the best-fit including all backgrounds. The blue curves are the expectation accounting for the distance to the individual reactors, time dependent flux variations and efficiencies. The error bar shows the statistical uncertainty only.

References

- [1] K. Eguchi *et al.*, Phys. Rev. Lett. **90**, 021802 (2003).
- [2] T. Araki *et al.*, Phys. Rev. Lett. **94**, 081801 (2005).
- [3] T. Araki *et al.*, Nature **436**, 499 (2005).
- [4] S. Abe *et al.*, Phys. Rev. Lett. **100**, 221803 (2008).
- [5] S. Harissopulos, H. W. Becker, J. W. Hammer, A. Lagoyannis, C. Rolfs and F. Strieder, Phys. Rev. C **72**, 062801 (2005).
- [6] JENDL 2005 <http://www.ndc.tokai-sc.jaea.go.jp/jendl/jendl.html>
- [7] S. Agostinelli *et al.*, Nucl. Instrum. Meth. A **506**, 250 (2003).